

E L A T E. *Gen. pl.* 1226.

MASC. *Cal.* 3-dentatus. *Cor.* 3-petala. *Antheræ*
6, sessiles.

FEM. *Cal.* 1-phyllus. *Cor.* 3-petala. *Pist.* 1. *Stigmata* 3. *Drupe* 1-sperma.

1. *E.* frondibus pinnatis: foliis oppositis. *Sp. pl.* 1659. *Sylvestris.*
Prickly-leav'd Elate.

Nat. of the East Indies.

Intro. about 1763, by John Blackburne, Esq.

Fl. S. F.

Z A M I A. *Gen. pl.* 1227.

MASC. *Amentum* strobiliiforme, squamis subtus tectis polline.

FEM. *Amentum* strobiliiforme, squamis utroque margine. *Bacca* solitaria.

1. *Z.* foliis cuneiformibus rectis glaberrimis a medio ad apicem ferratis subtus furfuraceis, stipite spinoso. *furfuracea.*
Linn. fl.

Palma americana foliis Polygonati brevioribus leviter ferratis et nonnihil spinosis, trunco crasso. *Pluk. alm.* 276. t. 103. f. 2. et 309. f. 5.

Palma americana crassis rigidisque foliis. *Herm. parad.* 210. t. 210.

Palmifolia femina. *Trew. chret.* 5. t. 26.

Broad-leav'd Zamia.

Nat. of the West Indies.

Cult. 1691, in the Royal Garden at Hampton-court.

Pluk. loc. cit.

Fl. July and August. S. F.

*integri-
folia.*

- 2.
- Z. foliolis subintegerrimis obtusiusculis muticis rectis nitidis, stipite inermi. Linn. fl.*

*Zamia pumila. Sp. pl. 1659. (exclusis synonymis.)*Dwarf *Zamia*.*Nat. of East Florida.**Introd. 1768, by John Ellis, Esq.**Fl. July and August.*

S. ƒ.

debilis.

- 3.
- Z. foliolis linearibus muticis apice ferrulatis patenti-recurvis rachi canaliculata longioribus, stipite tri-quetro compresso inermi. Linn. fl.*

Palma prunifera humilis non spinosa Insulæ Hispaniolæ, fructui jujubino similis, officulo triangulo. *Comm. hort. 1. p. 111. t. 58.*Long-leav'd *Zamia*.*Nat. of the West Indies.**Introd. 1777, by Messrs. Kennedy and Lee.**Fl. July and August.*

S. ƒ.

pungens.

- 4.
- Z. foliolis subulatis patentibus strictis rigidis mucronatis: margine exteriori baseos rotundato, stipite teretiusculo inermi. Linn. fl.*

Palma sobolifera, ægyptia, foliis levioribus, fructu nigro. *Till. pif. 129. tab. 45.*Needle *Zamia*.*Nat. of the Cape of Good Hope. Mr. Fr. Masson.**Introd. 1775.**Fl.*

G. H. ƒ.

Cycadis.

- 5.
- Z. foliolis obliquis lineari-lanceolatis subulatis pilosis curvatis apice uni-vel trispinosis, stipite inermi. Linn. fl.*

• *Zamia Cycadis. Linn. suppl. 443.**Cycas caffra. Thunb. in nov. act. upsal. 2. p. 283. t. 5.*

α foliis apice unispinosis.

Entire narrow-leav'd *Zamia*.

β foliis

β foliis apice bi-vel tri-spinosis.

Trifid narrow-leav'd *Zamia*.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson*.

Introd. 1775.

Fl.

G. H. 4.

A P P E N D I X.

γ. GINGKO. *Linn. mant.* 313. *Thunb. japon.* 358. *bioba.*
Maiden-hair Tree.

Nat. of Japan.

Cult. 1758, by Mr. James Gordon.

Fl.

H. 5.

ADDENDA,